SEDIMENT DISTRIBUTION AND COMPOSITION ON THE SHALLOW WATER CARBONATE BASIN OF THE ZANZIBAR CHANNEL

YW Shaghude

Institute of Marine Sciences, University of Dar es Salaam P.O. Box 668 Zanzibar

Received 16 September 2002: Accepted November 2003

ABSTRACT

Sediments of the shallow water carbonate basin in Zanzibar channel were investigated for composition and grain size distribution. The surface sediment composition was dominated by carbonate sands (with CaCO₃ > 30%), except in the area adjacent to mainland coastline and a thin lobe which projects from Ruvu River to the middle of the channel. Mean grain size distribution closely resembles that of the carbonate content, where Tidally Dominated Reef Platform Sediments (TDRPS) located east of the Zanzibar channel were characterised by medium to coarse sands and the siliciclastic sediments adjacent to the mainland were characterised by fine sand. The TDRPS are the most poorly sorted sediments with sorting values between 1.2 and 1.6 phi. The present study highlights existence of major differences between the eastern and western sides of the channel. Sediments on the eastern side of the channel, that are predominantly biogenic, were characterized by grain size frequency curves without any prominent mode. Sediments on the western side of the channel are composed of both biogenic and terrigenous material. The grain size frequency curves of these sediments have a fine mode and usually a coarse tail.

INTRODUCTION

In many parts of the world, shallow water marine environments are areas of extensive carbonate production and accumulation. Shallow carbonate deposits widely reported in warm shallow water shelves (e.g. Harris et al. 1990, 1996, Li et al. 1998) were viewed as tropical or sub-tropical phenomena. However, from the late 1960's there has been a dramatic increase in information regarding shallow carbonate deposits outside the tropical realm. Modern shallow carbonate deposits are now widely reported in the temperate shelves (Nelson, 1978, Nelson et al.1982, Nelson & Bornhold 1983, Carey et al. 199, Barnhardt & Kelley, 1995) as well as high latitudes

(Alexandersson 1979, Boreen et al. 1993, Taviani et al. 1993).

In many shallow water carbonate basins, the input of terrigenous sediments (to the basin) is one of the most important limiting factors, affecting both the carbonate production and accumulation. Studies in many parts of the world show that, as long as the input of terrigenous material is minimal, the potential exists for carbonate production and accumulation (Nelson & Bornhold 1983). Shallow marine carbonate producers use seawater to build their skeletons. The shallow marine carbonate producers include both the autotrophic primary producers (e.g. coralline algae) and other non-autotrophic benthic sessile organisms (e.g. molluscs, foraminifera,

echinoderms and sponges) as well as sedentary organisms such as corals or bryozoans. The turbidity associated with fluvial discharge of terrigenous sediment can reduce light penetration thereby reducing primary productivity. This would therefore prohibit shallow marine carbonate production by the autotrophic carbonate producers. Further, as most shallow water benthic carbonate producers are suspension feeders, the fine terrigenous materials may clog the body feeding system of these organisms (Carey et al. 1995) and thus inhibit their growth. Finally, high input of terrigenous material may bury the benthic carbonate producing organisms and effectively reduce the size of the carbonate basin.

The present study investigated some of the important characteristics of the sediments on shallow water carbonate basin of the Zanzibar channel (Fig. 1-a), both in terms of sediment composition and grain size distribution. Further investigations were made on how terrigenous sediments are redistributed in the basin. A section of the Zanzibar channel between the two major rivers (the Ruvu and Wami), which is approximately one third of the channel was covered (Fig. 1-a).




Fig 1: Map of coastal Tanzania showing (a) approximate boundary of the investigated area and (b) the grab sampling. Note the high density data in the Tidally Dominated Reef Platform Sedients (TDPRS). The reef platforms are shown as non-hatched polygons and the hatched polygons in the central and western sides of the Zanzibar channel are patched reefs.

METHODS

Sediment samples were collected using a light (approx. 10 kg) Van Veen grab sampler, with a carrying capacity of 2 to 3 kg. The samples were collected in two phases (Fig. 1-b); the first phase, characterised by dense sampling sites (100 samples) close to Zanzibar town and the second phase characterised by less dense sampling sites (40 samples) extended across the channel.

The sediment samples were washed with distilled water and then oven-dried at 45°C. All samples were analysed for grain size distribution by dry sieving through -1 phi (2 mm) and 4 phi (63 µm) sieves, with successive sieves spaced at ý phi intervals. Grain sizes for the sieved sediments were evaluated by calculating the graphic mean, sorting, skewness and kurtosis (Folk & Ward 1957). Carbonate content in the sediments was determined by the leaching method using dilute hydrochloric acid Approximately 2 g of well-ground subsamples were leached and the carbonate content calculated from weight loss during leaching.

RESULTS

Carbonate sediments (>30% CaCO₃) occupied more than 70% of the sea bottom (Fig. 2-a), with the highest carbonate content (>80%) in the Tidally Dominated Reef Platform Sediments (TDRPS) on the eastern side of the channel (Shaghude *et al.* 2001). Siliciclastic sediments (< 30% CaCO₃) dominated the coastal strip about 5 km adjacent to the mainland except at river Ruvu where a lobe of siliciclastic sediments projects from the river mouth to nearly the middle of the channel.

The siliciclastic sediments were generally very minor components in the TDRPS, confined on the intertidal zone adjacent to Zanzibar Island. The remaining part of the sea bottom may be described as an intermediate carbonate zone (30%<CaCO₃<80%).

The distribution of mean grain size (Fig. 2-b) in the general perspective closely fits the distribution of the carbonate content. In particular, the TDRPS which are characterised by the highest carbonate content, were of medium (1.0 to 2.0 phi) to coarse (1.0 to 0.0 phi) sand size, while the siliciclastic dominated zone adjacent to the mainland coast was characterised by fine (2.0 to 3.0 phi) sand sediments.

The TDRPS (Fig. 2-c) were the most poorly sorted sediments with sorting values between 1.2 and 1.6 phi. On the Folk and Ward (1957) sorting scale; these sediments may be described as poorly sorted. The poor sorting value in these sediments is probably due to the fact that they have various kinds of biogenic components namely benthic foraminifera, pelecypods, gastropods, corals and ostracods (Shaghude & Wannäs 2000). The best-sorted sediments in the channel were the siliciclastic ones close to the mainland coastline, with sorting value less than 1.2 phi. The relatively better sorting nature of these sediments, compared to other sediments, is probably due to their fine sized nature (dominated by sediments of siliciclastic origin); fine sediments are often better sorted than coarse sediments (Folk, 1964). The sediments in the central parts of the channel were also poorly sorted but their sorting range was lower than that of reef platform sediments. The sorting value of most of these sediments was close to 1.2 phi.




Fig. 2: Map showing (a) the distribution of carbonate content (b) the main grain size distribution, (c) sorting, (d) skewness and (e) kurtosis of the investigated area, Zanzibar Channel. The small cross symbols in the figure are the sampling points. Note the siliclastic/ carbonate sediments boundary (a) shown by the dotted lines (30% CaCO₃).

Skewness of the sediments (Fig. 2-d) varied from positive skewness (about 0.2) to being very negatively skewed (about -0.5). Investigation of the cross-shore skewness trend shows that the skewness generally changes from symmetrical or positive skewness near the shore to negatively or very negatively skewed sediments at the middle of the channel. The highest peakedness (kurtosis) in the sediments (Fig. 2-e) was in the southern parts of the investigated area, with peakedness varying between 0.8 and 1.6. The sediments on the central and northern parts had peakedness values between 0.8 and 1.2. Although further attempt may be made to interpret these values, due to low-density data in the western and southern parts of the study area, interpretation based on skewness and kurtosis, which are third and fourth statistics, respectively, is considered to be unnecessary.

DISCUSSION

To a large extent the two rivers Ruvu and Wami on the mainland control the sea bottom morphology and the sediment composition on the western side of the channel. The two rivers are the major source of siliciclastic sediments on the western side of the channel. A major contribution of the sediments from the rivers Ruvu and Wami comes from the Uluguru and Nguru mountains which are tropical mountains located about 200 km from the coastline of the Tanzania mainland (Fig. 1-a). The bedrock geology of these mountains consists of metamorphic crystalline rocks of the Mozambique belt, believed to have been formed during the Pan African episode, about 550±100 m.y. b.p (Windley 1986), and probably uplifted before the Upper Carboniferous (Samson & Wright 1964). The local relief of the mountains is over 2600 m above sea level and the mountains generally receive high precipitation of over 1400 mm/ per annum (Temple & Rapp 1972, Rapp et al. 1972).

Topography (relief) and rainfall are generally considered to be the most important factors, controlling sediment discharge to the sea for most rivers (Milliman and Syvitski 1992, Milliman 1995). Thus, the overall high precipitation on these mountains and the high relief of the mountains are probably the most important factors which control the siliciclastic sediment discharge to the sea via the two rivers, Ruvu and Wami.

The Zanzibar channel is fairly protected from the effect of long period waves coming from the open ocean as it does not have a direct connection to the open ocean (Brampton 1996). However, the effects of locally generated short period waves cannot be neglected. Furthermore, as the tides are mesotidal (Boothroyd 1985), they are considered to be moderate. Both tides and waves are therefore considered to control the redistribution of the sediment laden river output from the river mouths to other parts of the coastal area on the Tanzania mainland. Exceptional development of beach ridges immediately north of the river mouths on the Tanzania mainland has been pointed out as an evidence of preferential northward transport of sediments along the shore (Alexander 1969). Thus, it appears that most of the terrigenous materials are preferentially transported northward by the longshore drift. Exception to this may be at the river mouths or tidal channels where the ebb currents may carry the fine material further offshore as demonstrated by the observed siliciclastic lobe at river Ruvu.

The study of Shaghude and Wannäs (2000), who investigated the same study area discussed the mineralogical and biogenic composition of the sediments in the Zanzibar channel. This study proposed that the Zanzibar channel be divided into three bio-physiographic zones: 1-The coastal zone (0-10 m water depth), 2 –

the reef platforms/patch reefs zone (10-20 m) and 3 – the central channel zone (>20 m). This morphological division was supported by the results of the statistical analyses on the biogenic and mineralogical data as well as the spatial distribution of biogenic:quartz ratio. In particular, the statistical analyses showed significant differences in the spatial distribution of quartz, biogenic content and the biogenic to quartz ratio (for the thin sections analysed) among the three bio-physiographic zones.

While the three-fold depth division defining the bio-physiographic zones in the channel suggest existence of some general conclusions about the sediment characteristics, it does not highlight on the important difference between the eastern and western sides of the channel. This difference can be evidenced from the present results on grain size distribution parameters (namely the mean grain size and sorting), distribution of carbonate content and the sea bottom morphology. Figure 3 showing the grain size frequency curves from the 49 samples used in the study of Shaghude and Wannäs (2000) further highlights the difference between the eastern and western side of the channel. In the two-fold geographic division, the sediments east of the 30 m dashed contour (Fig. 3-e), that are essentially the TDPRS, are considered to be one division while the sediments west of this contour are considered to be another division.

The grain size frequency curves for the investigated samples (Fig. 3) indicate different depositional environments between the eastern and the western sides of the channel. The samples on the eastern side (Fig. 3-a) generally did not have a significant proportion of fine grain material. Instead, most of these samples contained coarse material of biogenic origin. and the samples did not have a prominent mode. Very exceptional cases (Fig. 3-b) include one near shore sample (No. 20) and the other sample near the central deep (No. 12). Both of these two samples had significant fine material similar to most of the samples on the western division (Fig.3-c). The samples on the western division had a high proportion of fine material and often had a coarse tail. The fine mode in these samples is not characteristic, but generally lies between 2 and 3.5 phi (0.25 and 0.09 mm). The extremely exceptional cases in the western division are the samples shown in Fig. 3-d. The grain size curves of these samples closely resembled those of the samples from the eastern division, possibly due to their proximity to reef platforms or patch reefs.




Fig. 3: Grain size distribution curves of the investigated samples. a - b = eastern side; c - d = western side; e = location of the 49 samples whose grain size distribution curves are discussed. Note that a and c represent the typical cases, while b and d represent the abnormal cases.

The present study highlights the major differences between the eastern and western side of the channel. The sea bottom on the eastern side of the channel was dominated by raised reef platforms with a network of internal channels between these platforms. Only a few patch reefs were evident on the western side of the channel, which was generally characterized by a smooth topography. The bulk of the sediments on the eastern side of the channel are of biogenic origin with very little contribution from the terrigenous source. The carbonate content in these sediments generally exceeded 60%. These sediments were of medium to course texture and poorly sorted, with sorting value > 1.2 phi. The grain size frequency curves of these sediments do not have any prominent mode. The sediments on the western side of the channel were composed of both biogenic and terrigenous material and the carbonate content in these sediments was generally less than 60%. These sediments were of fine to medium texture and had a better sorting value than the tdrps east of the channel. The sorting value in these sediments did not exceed 1.2. The grain size frequency curves of these sediments exhibited a fine mode and usually a coarse tail.

ACKNOWLEDGEMENT

The author gratefully acknowledges the financial assistance received from Sida/SAREC to run this project. I highly appreciate the guidence provided by Drs Kjell O. Wannäs and Eve Arnold from Stockholm University. The manuscript also benefited greatly from the comments of two anonymous reviewers.

REFERENCES

Alexander CS 1969 Beach ridges in northeastern Tanzania. Geographical review 59: 104-123

- Alexandersson ET 1979 Marine maceration of skeletal carbonates in Skagerrak, North Sea. *Sedimentology* **26**: 845-852
- Barnhardt WA and Kelley JT 1995 Carbonate accumulation on the inner continental shelf of Maine: A modern consequence of late Quaternary glaciation and sea level change. *Journal of Sedimentary Research*, **A65(1)**: 195-207
- Boreen T, James N, Wilson C and Heggie D 1993 Surficial cool-water carbonate sediments on the Otway continental margin, Southeastern Australia. *Marine* Geology 112: 35-56
- Boothroyd JC 1985 Tidal inlets and tidal deltas. In: (RA Davis, Jr. (eds) *Coastal sedimentary environments*. Springerverlag, pp.445-532.
- Brampton A 1996 Processes of beach change. In S.K. Mohammed and J. Betlem (eds) Proceedings of the National Workshop on coastal erosion Zanzibar, 12 April 1996. Zanzibar environmental study series, No. 20.
- Carey JS, Moslow TF and Barrie JV 1995 Origin and distribution of Holocene temperate carbonates, Hercate Strait, Western Canada Continental shelf. *Journal* of sedimentary Research A65(1): 185-194
- Folk RL and Ward WC 1957 Brazo River bar, a study in the significance of grain size parameters. *Journal of Sedimentary Petrology* **27**: 3-27
- Folk RL 1964 Carbonate sediments of Isla Perez Alakran Reef Complex, Yucatan. *Journal of Geology*, **72**: 255-292
- Harris PT, Davies PJ and Marshall JF 1990 Late Quaternary sedimentation on the Great Barrier Reef continental shelf and slope east of Townsville, Australia. *Marine* Geology 94: 55-77
- Harris PT, Pattiaratchi CB, Keene JB, Dalrymple RW, Gardener JV, Baker EK Cole AR, Mitchell, D Gibbs P and Schroeder WW 1996 Late Quaternary deltaic and carbonate sedimentation in the

- Harris PT, Pattiaratchi CB, Keene JB, Dalrymple RW, Gardener JV, Baker EK Cole AR, Mitchell, D Gibbs P and Schroeder WW 1996 Late Quaternary deltaic and carbonate sedimentation in the Gulf of Papua foreland basin: response to sea-level change. *Journal of Sedimentary Research* 66(4): 801-819
- Li C Jones B and Kalbfleisch WBC 1998 Carbonates sediment transport pathways based on foraminifera: Case study from Frank Sound, Grand Cayman, British West Indies. Sedimentology, 45: 109-120.
- Milliman JD 1995 Sediment discharge to the ocean from small mountainous rivers: The New Guinea example. *Geo-Marine Letters* **15**: 127-133
- Milliman JD and Syvitski JPM 1992 Geomorphic/tectonic control of sediment discharge to the ocean: the importance of small mountainous rivers. *Journal of Geology* 100: 525-544
- Nelson CS 1978 Temperate shelf carbonates sediments in the Cenozoic of New Zealand. Sedimentology 25: 737-771
- Nelson CS, Hancock GE and Kamp PJJ 1982 Shelf to basin, temperate skeletal carbonate sediments, Three Kings, New Zealand. *Journal of Sedimentary Petrology* **52**: 717-732
- Nelson CS and Bornhold BD 1983 Temperate skeletal carbonate sediments on Scott shelf, Northwestern Vancouver Island, Canada. *Marine Geology* **52**: 241-266

- Rapp A, Axelsson V Berry L and Hammond M 1972 Soil erosion and sediment transport in the Morogoro river catchment, Tanzania. *Geografiska Annaler*, **54A**: 125-155
- Samson DN and Wright AE 1964 The geology of the Uluguru mountains. *Bulletin of the geological survey of Tanganyika* 37
- Shaghude YW and Wannäs KO 2000 Mineralogic and biogenic composition of the Zanzibar channel sediments, Tanzania. Estuarine coastal and shelf science 51: 477-487
- Shaghude YW, Wannäs KO and Mahongo SB 2001 Hydrodynamic setting of the tidal dominated platforms and reef environment sediments of the Zanzibar channel. Presented during the second Western Indian Ocean Marine Science Association (WIOMSA) symposium, 22-26 October, 2001, Dar es Salaam
- Taviani MT, Reid DE and Anderson JB 1993 Skeletal and isotopic composition and paleoclimatic significance of Late Pleistocene carbonates, Ross Sea, Antarctica. Journal of sedimentary petrology 63(1): 84-90
- Temple PH and Rapp A 1972. Landslides in the Mgeta area, western Uluguru mountains, Tanzania. Geografiska Anneler 54A: 157-193
- Windley BF 1986 The evolving continents, Bass Press Avon, Britain